

Search for exotic transitions of muon neutrinos to electron
neutrinos with MINOS

MARIANNA GABRIELIAN

(for the MINOS and MINOS+ Collaborations)

*Department of Physics
University of Minnesota, Minneapolis, MN, USA*

The observed neutrino flavor transitions are currently explained by the three flavor neutrino oscillation phenomenon, considered to be the leading order mechanism behind the flavor transitions. Currently existing data from LSND, MiniBooNE and reactor experiments demonstrate anomalies that could potentially be indications of non-standard neutrino phenomena. MINOS can probe transitions from muon neutrinos to electron neutrinos and search for anomalous behavior that cannot be explained by standard model neutrino oscillations. Here we present the search for second order effects in the flavor transitions by analyzing the MINOS $\nu_\mu \rightarrow \nu_e$ channel.

PRESENTED AT

DPF 2015

The Meeting of the American Physical Society
Division of Particles and Fields
Ann Arbor, Michigan, August 4–8, 2015

1 Introduction

One of the motivations for exploring the sterile neutrino oscillations in the $\nu_\mu \rightarrow \nu_e$ appearance channel is the existing tension between the short (SBL) and long (LBL) baseline neutrino experiments. For example, the Liquid Scintillator Neutrino Detector (LSND) and MiniBooNE experiments report anomalous excess of $\bar{\nu}_e$ appearance in a $\bar{\nu}_\mu$ beam over a short baseline, which cannot be explained by standard three flavor oscillations [1, 2].




Figure 1: LSND and MiniBooNe allowed regions for the sterile mixing angle and mass splitting. Shows tension with the OPERA and ICARUS long baseline experiments.

One of the explanations is an existence of one or more sterile neutrino states that are not interacting via the weak interaction. Adding a new neutrino type to 3ν model introduces another mass splitting and three mixing angles to parameter space. Figure 1 shows the allowed regions for the additional mass splitting and $\theta_{\mu e}$ mixing angle. MINOS+ ν_e appearance analysis uses $(3+1)\nu$ model to investigate the sterile oscillation hypothesis in the parameter space allowed by LSND and MiniBooNE.

2 MINOS Experiment

The Main Injector Neutrino Oscillation Search (MINOS) experiment [3] is a long-baseline neutrino oscillation experiment utilizing the Neutrinos at the Main Injector (NuMI) neutrino beam. The experiment consists of two functionally identical tracking calorimeters, primarily optimized to detect μ^+/μ^- , separated by 735 km distance.

The main goals of the experiment are to make precise measurements of the neutrino oscillation parameters to other active neutrino flavors, explore the possibility of sterile oscillations as well as to study exotic scenarios such as non-standard neutrino interactions. Near Detector (ND) is located at Fermilab ~ 1 km away from NuMI target, is measuring the neutrino flux, cross section, energy spectrum and the flavor composition of the beam. Far Detector (FD) is located at Soudan Underground Laboratory at Northern Minnesota, ~ 735 km away from NuMI target, measures the same beam parameters. Functionally identical design was chosen to effectively cancel out systematic uncertainties related to the neutrino beam flux and neutrino interaction cross sections. Since the MINOS detectors were not optimized for ν_e CC




Figure 2: Simulated NuMI neutrino energy spectrum for MINOS, MINOS+ and NOvA experiments.

events, a new statistical selection algorithm was implemented to distinguish $\nu_\mu \rightarrow \nu_e$ candidates from background interactions - particularly NC events. ν_e appearance provides sensitivity to θ_{13} , θ_{23} octant, mass hierarchy and δ_{CP} .

MINOS+ is a continuation of the MINOS experiment, operating the same detectors. It will run concurrently with the NOvA experiment with an upgraded NuMI beam. MINOS+ allows access to higher energy region to study sterile neutrino oscillations as well as non-standard neutrino interactions, with beam covering wide range of energies and peaking at 7 GeV as can be seen from Figure 2.

3 NuMI Beamline

The ν_μ neutrino beam is produced by colliding 120 GeV proton beam from the Main Injector with graphite target. Mesons ($\pi^\pm/K^\pm$), produced as a result of these collisions, are then focused into the decay pipe by focusing magnetic horns, where they are




Figure 3: Schematic view of NuMI production beamline

allowed to decay (Figure 3). This admixture then passes through an absorber, which stops all charged particles from moving forward. As a result wide-band on-axis ν_μ beam is produced and directed towards the MINOS detectors. By switching the magnetic horn polarity, one can switch from neutrino to antineutrino beam production mode.

4 Sterile Neutrino Sensitivity in ν_e Appearance

The signature of $\nu_\mu \rightarrow \nu_s$ oscillations in MINOS manifests itself as an excess of ν_e -CC events in the Far Detector, depending upon the particular Δm_{41}^2 and neutrino energy at the same time causing energy-dependent depletion in both NC and ν_μ -CC energy spectra. This is demonstrated in Figure 4 for the shown parameters. Signal and Background Monte Carlo distributions show the excess in the signal and depletion in the background energy spectra in the presence of nonzero θ_{14} and θ_{24} mixing angles.

The sensitivity plots were generated by applying Gaussian intervals to the log-likelihood surfaces produced for specified values of Δm_{41}^2 . At each Δm_{41}^2 value, a new surface was produced by profiling the combinations of θ_{34} , δ_{CP} , and δ_{eff} . Figure 5 shows the 90% confidence level contour for MINOS+ in the SBL parameter space. The shaded blue region is excluded MINOS+.

The MINOS+ experiment is capable of decoupling the θ_{14} and θ_{24} . In Figure 6 the shaded regions correspond to LSND allowed regions. The blue dashed curves show the $\Delta m_{41}^2 = 0.15 \text{ eV}^2$ LSND region, while the red dashed lines outline the $\Delta m_{41}^2 = 0.50 \text{ eV}^2$ LSND region. The solid lines show the MINOS+ 90% C.L. contours for the same two values of Δm_{41}^2 . MINOS+ excludes the parameter space to the upper-right of the solid contours. For $\Delta m_{41}^2 = 0.15 \text{ eV}^2$ MINOS+ can exclude most of the LSND signal region, and although for $\Delta m_{41}^2 = 0.50 \text{ eV}^2$ MINOS's sensitivity is relatively poor, it still can exclude some portions of the LSND signal parameter space. Higher values of $\Delta m_{41}^2 (1 - 10^2 \text{ eV}^2)$ can also be probed, but this will require incorporating oscillations in the Near Detector when extrapolating to Far Detector.




Figure 4: Monte Carlo Signal and Background reconstructed energy distributions. Solid lines correspond to 3ν model predictions, while the dashed lines correspond $(3+1)\nu$ model predictions for the given values of parameters.




Figure 5: 90% confidence level contour shows the MINOS+ allowed region in the SBL parameter space, assuming both $\theta_{14}, \theta_{24} < 0.25\pi$.




Figure 6: 90% C.L. in two-dimensional parameter space for two chosen values of Δm_{41}^2 , assuming $\theta_{14}, \theta_{24} < 0.25\pi$. MINOS+ excludes the parameter space to the upper-right of the solid contours. The shaded bands represent the LSND signal region associated with the indicated values of Δm_{41}^2 .

5 Summary

In conclusion, MINOS+ is collecting new high-statistics data with medium energy beam. This will allow new precision measurements of 3-flavor oscillation parameters in an unexplored energy range. It opens pathway to probe sterile neutrino oscillation search in ν_e appearance channel as well as explore non-standard neutrino interaction scenarios in the 6-12 GeV region. MINOS+ ν_e appearance can be used to set new constraints on low-mass sterile mixing parameters.

ACKNOWLEDGMENTS

This work was supported by the U.S. DOE; the United Kingdom STFC; the U.S. NSF; the State and University of Minnesota; Brazil's FAPESP, CNPq and CAPES. We are grateful to the Minnesota Department of Natural Resources and the personnel of the Soudan Laboratory and Fermilab for their contributions to the experiment.

References

- [1] LSND Collaboration: A.Aguilar et al., Evidence for neutrino oscillations from the observation of ν_e appearance in a ν_μ beam, Phys. Rev. **D64**, 112007 (2001).
- [2] MiniBooNE Collaboration: A.A. Aguilar-Arevalo et al., Improved Search for $\nu_\mu \rightarrow \nu_e$ Oscillations in the MiniBooNE Experiment, Phys. Rev. Lett. **110**, 161801 (2013).
- [3] MINOS Collaboration: D.G Michael et al., The magnetized steel and scintillator calorimeters of the MINOS experiment, Nucl. Instrum. Meth. **A 596**, 190 (2008).